



# The Disenchantment in Modern Physics

Wim Vegt 

1. Department of Physics, Eindhoven University of Technology, The Netherlands

## Abstract:

Isaac Newton looms over the Enlightenment as the giant who steered humanity out from under the dark shadows of superstition and into the bright, sterile light of the mechanical universe. Armed with the navigational tools of empirical evidence and mathematical proofs, he deciphered the forces of nature using human reason alone. With the publication of his masterwork *Principia* in 1687, the archaic belief in Nature's soul was left shipwrecked on the shores of the unscientific past. Newton's legacy is scientific positivism, or the belief that an idea is true only when it can be proved mathematically. If, in collective memory, René Descartes (1596-1650) succeeded in splitting the mind from the body, then Newton's clockwork universe desecralized nature and cleaved the body from the soul. [43] At the end of 1918, Max Weber proclaimed at the University of Munich that 'the fate of our times' is the 'disenchantment of the world' The understanding that mystery and sacredness was disappearing from a world increasingly dominated by industry, science, and technology had its intellectual roots in German Romanticism, where it had found expression in the works of Novalis and Friederich Hölderlin. After Weber reintroduced the evocative term, "Entzauberung", it would go on to have a deep impact on theories of modernity developed by the following generations.[44] Newton dressed his mystical intuition in mathematical proofs before presenting his ideas to the world. The father of mechanical science believed that spirit animated matter and his work was to understand, using the gifts of his meager human intellect, the eternal and divine. This article focusses on the "Prima Materia" described by Isaac Newton [43] and already thousands of years ago in "The Emerald Tablet of Hermes Trismegistus". [45] In the Emerald Tablet, the "One Thing" refers to the "Prima Materia". This was then acted upon by the thoughts or word of the One Mind to create the material reality we can observe with our senses. The fundamental concept of the "Prima Materia" has been expressed in Equation (16) and (17) in which "Light" is the "Prima Materia" of our reality and the origin of our universe. Light is the building material out of which this universe has been created. The new theory, describing electro-magnetic-gravitational-acceleration force density interactions (expressed in  $N/m^3$ ), has been discussed at astronomical levels: Gravitational RedShift, Black Holes and Dark Matter and at sub-atomic levels: The absorption and emission of light at sub-atomic levels in concentric spheres by an atom at discrete energy levels. Evidence will be demonstrated about the correctness of this new electro-dynamic theory which represents the only theory which connects electro-dynamics in a correct way with plasma-dynamics. In general, the gravitational (acceleration) force densities, originating from rotation and linear accelerations, are being ignored but with nuclear fusion processes these gravitational (acceleration) forces become fundamental and are necessary to develop a stable nuclear fusion process. Differently than in General Relativity, the electro-magnetic-gravitational-acceleration force density interactions (expressed in  $N/m^3$ ) [35] fundamentally has been based on the divergence of the sum of the "Stress Energy Tensor" and the introduced "Gravitational-Acceleration" Tensor. The theory describes "Gravitational-Acceleration-Electromagnetic" Interaction resulting in a mathematical Tensor presentation for BLACK HOLES. (Gravitational Electromagnetic Confinements) [1] The

“Electromagnetic Energy Gradient” creates a second order effect “Lorentz Transformation” which results in the Gravitational Field of BLACK HOLES which determines the interaction force density between the confinement of Light (BLACK HOLE’s) and the “Gravitational-Acceleration” Field. Einstein approached the interaction between gravity and light by the introduction of the “Einstein Gravitational Constant” in the 4-dimensional Energy-Stress Tensor  $\kappa T_{\mu\nu}$  (1) In this alternative approach related to General Relativity, the interaction between gravity and light has been presented by the sum of the Electromagnetic Tensor  $T_{\mu\nu}$  and the “Gravitational-Acceleration” Tensor  $J_{\mu\nu}$  (2) The new theory describes the impact of "CURL" [38] within the gravitational fields around Black Holes and the impact on Gravitational Lensing. Gravitational "CURL" (Equation 6) is an effect which cannot be explained and calculated by General Relativity. The new approach presents mathematical solutions for the BLACK HOLES (Gravitational Electromagnetic Interaction) introduced in 1955 by Jonh Archibald Wheeler in the publication in Physical Review Letters in 1955 [1]. The mathematical solutions for BLACK HOLES are fundamental solutions for the relativistic quantum mechanical Dirac equation (Quantum Physics) in Tensor presentation (41). Assuming a constant speed of light “c” and Planck’s constant  $\hbar$  within the BLACK HOLE, the radius “R” of the BLACK HOLE with the energy of a proton, is about 1% of the radius of the hydrogen atom (14). The New Theory has been tested in an experiment with 2 Galileo Satellites and a Ground Station by measuring the Gravitational RedShift in a by the Ground Station emitted stable MASER frequency [2]. The difference between the calculation for Gravitational RedShift, within the Gravitational Field of the Earth, in “General Relativity” and the “New Theory” is smaller than  $10^{-16}$  (12) and (13). In all “General Redshift Experiments” “General Relativity” and the New Electrodynamical-Gravitational-Acceleration Theory predict a Gravitational RedShift with a difference smaller than 15 digits beyond the decimal point which is beyond the accuracy of modern “Gravitational Redshift” observations. Both values are always within the measured Gravitational RedShift in all observations being published since the first observation of the gravitational redshift in the spectral lines from the White Dwarf which was the measurement of the shift of the star Sirius B, the white dwarf companion to the star Sirius, by W.S. Adams in 1925 at Mt. Wilson Observatory. Theories which unify Quantum Physics and General Relativity [32], like “String Theory”, predict the non-constancy of natural constants. Accurate observations of the NASA Messenger [11] observe in time a value for the gravitational constant “G” which constrains until (  $|G|/G$  to be  $< 4 \times 10^{-14}$  per year). One of the characteristics of the New Theory is the “Constant Value” in time for the Gravitational Constant “G” in unifying General Relativity and Quantum Physics.

*Keywords: Quantum Physics, General Relativity, Gravitational RedShift, Black Holes, Dark Matter, Nuclear Fusion, Nuclear Plasmas.*

### AN ALTERNATIVE APPROACH IN GRAVITY AND ACCELERATION

Einstein approached the interaction between gravity and light by the introduction of the “Einstein Gravitational Constant” in the 4-dimensional Energy-Stress Tensor.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \quad (1)$$

In which  $G_{\mu\nu}$  equals the Einstein Tensor,  $g_{\mu\nu}$  equals the Metric Tensor,  $T_{\mu\nu}$  equals the Stress-Energy tensor,  $\Lambda$  equals the cosmological constant and  $\kappa$  equals the Einstein gravitational constant.

An alternative approach to Einstein's expression with the tensor  $\kappa T_{\mu\nu}$ , describing the curvature of the Space-Time continuum, is the sum of the Electromagnetic Tensor  $T_{\mu\nu}$  and the "Gravitational-Acceleration" Tensor  $J_{\mu\nu}$ .

$$\kappa T_{\mu\nu} \Leftrightarrow T_{\mu\nu} + J_{\mu\nu} \quad (2)$$

The 4-dimensional divergence of the sum of the Electromagnetic Stress-Energy tensor and the Gravitational Tensor expresses the 4-dimensional Force-Density vector (expressed in [N/m<sup>3</sup>] in the 3 spatial coordinates) as the result of Electro-Magnetic-Gravitational interaction.

$$f^\mu = \partial_\nu (T^{\mu\nu} + J^{\mu\nu}) \quad (3)$$

In vector notation the 4-dimensional Force-Density vector can be written as:

$$\vec{f}^4 = \begin{pmatrix} f_4 \\ f_3 \\ f_2 \\ f_1 \end{pmatrix} = \square \square (\vec{\bar{T}} + \vec{\bar{J}}) \quad (4)$$

The fundamental boundary condition for this alternative approach to gravity is the requirement that the Force 4 vector equals zero in the 4 dimensions, expressing a universal 4-dimensional equilibrium:

$$\vec{f}^4 = \begin{pmatrix} f_4 \\ f_3 \\ f_2 \\ f_1 \end{pmatrix} = \square \square (\vec{\bar{T}} + \vec{\bar{J}}) = \vec{0}^4 \quad (5)$$

The 3 spatial components of the Force-Density vector, as a result of Electro-Magnetic-Gravitational interaction can be written as:

$$\begin{aligned} \bar{f} = & -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \varepsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \varepsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \\ & + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \gamma_0 \bar{\mathbf{g}} (\nabla \cdot \bar{\mathbf{g}}) - \gamma_0 \bar{\mathbf{g}} \times (\nabla \times \bar{\mathbf{g}}) = \bar{0} \quad [\text{N/m}^3] \end{aligned}$$

in which:

$$\begin{aligned} \varepsilon_0 (\nabla \cdot \bar{\mathbf{E}}) &= \rho_E \text{ Electric Charge Density } [\text{C/m}^3] \\ \mu_0 (\nabla \cdot \bar{\mathbf{H}}) &= \rho_M \text{ Magnetic Flux Density } [\text{Vs/m}^3] \text{ or } [\text{Wb/m}^3] \\ \gamma_0 (\nabla \cdot \bar{\mathbf{g}}) &= \rho_M \text{ Mass Density (Electromagnetic) } [\text{kg/m}^3] \end{aligned} \quad (6)$$

Electric Energy Density:  $w_E = \frac{1}{2} \varepsilon_0 E^2$

Magnetic Energy Density:  $w_M = \frac{1}{2} \mu_0 H^2$

Gravitational Energy Density:  $w_G = \frac{1}{2} \gamma_0 g^2$

$\bar{\mathbf{g}}$  □ acceleration (gravitational or mechanical/dynamical)

In which E represents the electric field intensity expressed in [V/m], H represents the magnetic field intensity expressed in [A/m] and g represents the gravitational acceleration expressed in [m/s<sup>2</sup>]. The permittivity indicated as  $\varepsilon_0$ , the permeability indicated as  $\mu_0$  and the gravitational permeability of vacuum as  $\gamma_0$ .

For curl-free gravitational fields equation (6) can be written as:

$$\begin{aligned} \bar{f} = & -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \varepsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \varepsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \\ & + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \bar{\mathbf{g}} \rho_M = \bar{0} \quad [\text{N/m}^3] \end{aligned} \quad (7)$$

Substituting Einstein's  $W = m c^2$  in (7) results in "Electro-Magnetic-Gravitational Equilibrium Field Equation" (8):

$$\begin{aligned} \bar{f} = & -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \varepsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \varepsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \\ & + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \frac{1}{2c^2} \bar{\mathbf{g}} (\varepsilon E^2 + \mu H^2) = \bar{0} \quad [\text{N/m}^3] \end{aligned} \quad (8)$$

The theory describes "Electromagnetic-Gravitational Interaction", "Magnetic-Gravitational Interaction" and "Electric-Gravitational Interaction". In this new theory particles do not interact with fields. The interaction between an electric charged particle and an electric field is not the interaction between a particle and a field but it is the interaction between the electric field of the particle interacting with the other electric field. Every interaction is an interaction between fields. Electric Fields interact with Electric Fields, Magnetic Fields interact with Magnetic Fields and Gravitational Fields interact with Gravitational Fields.

## "GRAVITATIONAL REDSHIFT/ BLUESHIFT IN "LIGHT (EMR)" DUE TO "ELECTROMAGNETIC GRAVITATIONAL INTERACTION"

To test the New Theory, the Gravitational-Redshift experiment: "Test of the Gravitational Redshift with Galileo Satellites in an Eccentric Orbit" by S. Hermann et al., has been chosen [2]. In this experiment a stable "MASER" frequency from a ground station has been emitted to 2 Galileo Satellites, measuring the frequency difference between the Ground Station and the Satellites. The frequency shift has been caused by the gravitational field of the Earth and 2 satellites has been chosen to compensate for the eccentricity of the Galileo Orbit.

Assuming a gravitational field  $g[z]$  depending on the radial direction in cartesian coordinates between the ground station and the satellites:

$$\overline{g[z]} = \left\{ 0, 0, \frac{G M_{Earth}}{4 \pi z^2} \right\} \quad (9)$$

In which  $G$  ( $G = 6.67428 \cdot 10^{-11} \text{ Nm}^2 / \text{kg}^2$ ) equals the Gravitational constant,  $M_{Earth}$  the mass of the earth and  $r$  the radial distance from the centre of the earth. The mathematical solution [5] of equation (8) for plane electromagnetic waves (expressed in cartesian  $\{x,y,z\}$  coordinates) related to the Electric Field Intensity equals:

$$\overline{\mathbf{E}} = \begin{pmatrix} E_x \\ E_y \\ E_z \end{pmatrix} = \begin{pmatrix} e^{-\frac{G M_{Earth} \epsilon_0 \mu_0}{8 \pi z}} \hbar \left[ \omega_0 e^{-\frac{G M_{Earth} \epsilon_0 \mu_0}{4 \pi z}} (t - \sqrt{\epsilon \mu} z) \right] \\ 0 \\ 0 \end{pmatrix} \quad (10)$$

And the mathematical solution of (8) for the Magnetic Field Intensity equals:

$$\overline{\mathbf{H}} = \begin{pmatrix} H_x \\ H_y \\ H_z \end{pmatrix} = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{\epsilon_0 \mu_0}} e^{-\frac{G M_{Earth} \epsilon_0 \mu_0}{8 \pi z}} \hbar \left[ \omega_0 e^{-\frac{G M_{Earth} \epsilon_0 \mu_0}{4 \pi z}} (t - \sqrt{\epsilon \mu} z) \right] \\ 0 \end{pmatrix} \quad (11)$$

In which  $\omega_0$  equals the original frequency of the MASER radiation propagating in the direction of the gravitational field  $g[z]$  of the Earth in the  $z$ -direction. The exponential term demonstrates the Gravitational Redshift when the MASER radiation propagates in the direction of the Gravitational Field of the earth. The propagation speed of the Electromagnetic Radiation remains constant (the speed of light). But the amplitude of the field intensity and the frequency of the field intensity diminishes exponentially.

Calculations in Mathematica [5] demonstrate a difference between the calculation with General Relativity and the calculation with the New Theory. Choosing for the ground station a distance to the centre of the earth  $z_1 = 6,378,000$  [m] (Radius of the Earth) and for the average distance of the ESA satellites in a Galileo orbit  $z_2 = 23,222,000$  [m] (distance from the ESA satellite to the centre of the Earth), calculated with Mathematica, the Gravitational RedShift according General Relativity equals:

$$\Delta \omega_{GR} = 0.00000000004011815497097883 \text{ [s}^{-1}\text{]} \quad (12)$$

Calculated with Mathematica, the Gravitational RedShift according the New Theory, which is a solution of equation (8) equals:

$$\Delta \omega_{GR} = 0.00000000004011824206173742 \text{ [s}^{-1}\text{]} \quad (13)$$

Both calculated values a within the Range of the measured gravitational RedShift by the average values of both ESA satellites in the Galileo orbit

$$\Delta \omega_{Measured} = 0.000000000040118 \pm 2.2 \cdot 10^{-15} \text{ [s}^{-1}\text{]} \quad (14)$$

In [2] a factor  $\alpha$  has been defined which presents the measured deviation  $\alpha$  between the predicted Gravitational RedShift by General Relativity and the Measured Gravitational RedShift.

$$\alpha = \Delta \omega_{MEASURED} - \Delta \omega_{GR} = (2.2 \pm 1.6) \times 10^{-5} \quad (15)$$

A comparable factor  $\alpha$  can be used to determine which theory (General Relativity or the New Theory) has the nearest approach to the experimentally measured data. Highly accurate measuring experiments are required with an accuracy higher than 16 digits beyond the decimal point.

### BLACK HOLES

#### Black Holes without Singularities with Dimensions Smaller than the Diameter of the Hydrogen Atom

A second fundamental solution for equation (8) describes a Gravitational Electromagnetic Confinement (BLACK HOLE) [1] within a radial gravitational field with acceleration  $\bar{g}$  (in radial direction). This solution represents a Black Hole, the confinement of light due to its own gravitational field, and has no singularities. This solution for equation (8) describes Black Holes, dependent of time and radius, presenting discrete spherical energy levels, within a radial gravitational field with acceleration  $\bar{g}$  (in radial direction) [14] has been represented in (16) and (17).

$$\begin{pmatrix} E_r \\ E_\theta \\ E_\phi \end{pmatrix} = \begin{pmatrix} 0 \\ f(r) \sin(kr) \sin(\omega t) \\ -f(r) \cos(kr) \cos(\omega t) \end{pmatrix} \quad \begin{pmatrix} H_r \\ H_\theta \\ H_\phi \end{pmatrix} = \sqrt{\frac{\epsilon}{\mu}} \begin{pmatrix} 0 \\ -f(r) \sin(kr) \cos(\omega t) \\ -f(r) \cos(kr) \sin(\omega t) \end{pmatrix} \quad \bar{g} = \begin{pmatrix} \frac{G_1}{4\pi r^2} \\ 0 \\ 0 \end{pmatrix} \quad (16)$$

$$w_{em} = \left( \frac{\mu_0}{2} (\bar{m} \cdot \bar{m}) + \frac{\epsilon_0}{2} (\bar{e} \cdot \bar{e}) \right) =$$

$$f(r)^2 \left( (\sin(kr) \sin(\omega t))^2 + (\cos(kr) \cos(\omega t))^2 + \frac{\epsilon}{\mu} (\sin(kr) \cos(\omega t))^2 + (\cos(kr) \sin(\omega t))^2 \right)$$

In which the radial function  $f(r)$  equals:

$$f[r] = K e^{-\frac{G M_{BH} \epsilon_0 \mu_0}{8 \pi r}} \quad (17)$$

G represents the Gravitational constant and M represents the total confined electromagnetic mass of the BLACK HOLE. Equation (16) presents a Standing (Confined) Electromagnetic Field Configuration with a phase shift of 90 degrees between the electric field and the magnetic field with the corresponding Nodes and AntiNodes. [13]. The solution has been calculated according Newton's Shell Theorem, resulting in the fundamental concept of the "Prima Materia" [43] Assuming a constant speed of light "c" and Planck's constant ħ within the BLACK HOLE, the radius "R" (with n = 1,2,3,4....) of the BLACK HOLE with the energy of a proton, according W = m<sub>proton</sub> c<sup>2</sup>, would be: 1.5009211 × 10<sup>-10</sup> [J].

$$R_{GEON} = n \lambda = n \left( \frac{c}{f} \right) = n \left( \frac{c}{W} \right) \hbar = 7.1865 \cdot 10^{-26} \left( \frac{n}{W} \right) \quad (18)$$

$$R_{GEON} = n \cdot 3.82 \cdot 10^{-12} \text{ [m]}$$

Black Holes are varying from atomic dimensions with dimensions of 10<sup>-27</sup> [kg], Page 39 [33] until Black Holes with dimensions of 10<sup>40</sup> [kg], Page 67 [34]. At these dimensions Black Holes turn into Dark Matter. The fundamental boundary condition for the confinement of Electromagnetic radiation (BLACK HOLES) is that the energy flow (Poynting vector)  $\vec{S} = \vec{E} \times \vec{H}$  equals zero at the surface of the confinement. This is possible at every "90 degrees Phase Shift Surface" (Sphere) between the Electric Field and the Magnetic Field.

### Black Holes with a Singular Point and Large Dimensions

Fig 1 represents a Black Hole with a mass of 10<sup>35</sup> [kg] and a radius of about 25 [km] controlled by a different mathematical solution for equation (8). The radius of the Black Hole equals about 25 [km] which has been controlled by a different mathematical solution (19) for equation (8).

$$f[r] = K e^{\left( \frac{G M_{BH} \epsilon_0 \mu_0}{8 \pi r} - \log[r] \right)} \quad [\text{J} / \text{m}^3] \quad (19)$$





Fig. 1: The Energy Density [ J/ m<sup>3</sup>] as a function of the Radius R = max 10<sup>7</sup> [m] of the Black Hole.




**Fig. 2: The Energy Density [ J / m³ ] as a function of the Radius R = max 10<sup>5</sup> [m]**

Figure 1 and Figure 2 demonstrate the large effect of “Gravitational Intensity Shift” and “Gravitational RedShift” at the distance of 25 [km]. Over a distance of 10.000 [km] the intensity of the emitted light of the Black Hole with a mass of 10<sup>35</sup> [kg] falls back with a factor of 10<sup>-51</sup>. Also, the frequency of the emitted light of the Black Hole falls back with a factor 10<sup>-51</sup>. Emitted light in the visible spectrum of 10<sup>14</sup> [Hz] falls back to a frequency of 10<sup>-37</sup> [Hz]. These extreme low frequencies with extreme low intensities have never been measured which has result in the name “Black Hole” for the phenomenon of “Gravitational Intensity Shift” and “Gravitational RedShift” for a large mass. It follows from equation (8) and the solutions (10) and (11) that the speed of light does not change inside and around Black Hole. Only the direction of the propagation of light can change due to a gravitational field.

**Dark Matter in the Universe Controlled by “Gravitational Shielding”**


Fig 3 represents Dark Matter with a total mass of 10<sup>53</sup> [kg] and a radius of about 10 times the size of the Milky Way Galaxy. The radius of the dark mass equals 5 10<sup>21</sup> [m] which has been controlled by a different mathematical solution (20) for equation (8).

$$f [r] = K e^{\left( \frac{G M_{BH} \epsilon_0 \mu_0}{8 \pi r} - \log[r] \right)} \quad [J / m^3] \quad (20)$$



**Fig. 3: The Energy Density [ J / m³ ] as a function of the Radius R = max 10<sup>25</sup> [m] of the Dark Matter.**





**Fig. 4: The Energy Density [ J/ m<sup>3</sup>] of the Dark Matter as a function of the Radius R = max 10<sup>22</sup> [m]**

Figure 3 and Figure 4 demonstrate the large effect of “Gravitational Intensity Shift” and “Gravitational RedShift” at the distance of 5 10<sup>21</sup> [m] which is 10 times the radius of the Milky Way Galaxy. Over the distance of 5 10<sup>21</sup> [m] the intensity of the emitted light of the Dark Matter with a mass of 10<sup>53</sup> [kg] falls back with a factor of 10<sup>-261</sup>. Also, the frequency of the emitted light of the Black Hole falls back with a factor 10<sup>-261</sup>. Emitted light in the visible spectrum of 10<sup>14</sup> [Hz] falls back to a frequency of 10<sup>-247</sup> [Hz]. These extreme low frequencies with extreme low intensities have never been measured which has result in the name “Dark Matter” for the phenomenon of “Gravitational Intensity Shift” and “Gravitational RedShift” for an extreme large mass. It follows from equation (8) and the solutions (10) and (11) that the speed of light does not change inside and around the Dark Mass. Only the direction of the propagation of light can change due to the gravitational field of the Dark Mass.

### THE RELATIONSHIP BETWEEN BLACK HOLES AND QUANTUM PHYSICS

Introducing the Quantum Vector Function  $\bar{\phi}$  ,

$$\bar{\phi} = \sqrt{\frac{\mu}{2}} \left( \bar{H} + i \frac{\bar{E}}{c} \right) \tag{21}$$

Substituting (21) in (16) results in the quantum presentation for the BLACK HOLE:

$$\overline{\Phi(r,\theta,\varphi)} = \sqrt{\frac{\mu}{2}} \left( \bar{H} + i \frac{\bar{E}}{c} \right) f(r) \begin{pmatrix} \Phi_r \\ \Phi_\theta \\ \Phi_\varphi \end{pmatrix} \tag{22}$$

$$\overline{\Phi(r,\theta,\varphi)} = K \sqrt{\frac{\epsilon}{\mu}} e^{-\frac{G1 \epsilon_0 \mu_0}{8 \pi r}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\text{Sin}(k r) & \text{Sin}(k r) \\ 0 & -i \text{Cos}(k r) & i \text{Cos}(k r) \end{pmatrix} \begin{Bmatrix} 0 \\ \text{Cos}(\omega t) \\ i \text{Sin}(\omega t) \end{Bmatrix}$$

With "K" a constant value dependend of the mass of the BLACK HOLE. The Dot product between the unit vector and the Quantum Vector Function  $\bar{\phi}$  represents the quantum mechanical probability function  $\Psi[r, t]$  which is a fundamental solution of the Schrödinger Wave Equation.

$$\overline{\Phi(r, \theta, \varphi)} = K \sqrt{\frac{\epsilon}{\mu}} e^{-\frac{G1 \epsilon_0 \mu_0}{8 \pi r}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\text{Sin}(k r) & \text{Sin}(k r) \\ 0 & -i \text{Cos}(k r) & i \text{Cos}(k r) \end{pmatrix} \begin{Bmatrix} 0 \\ \text{Cos}(\omega t) \\ i \text{Sin}(\omega t) \end{Bmatrix} \quad (23)$$

$$\Psi(r, t) = \{1 \ 1 \ 1\} \begin{Bmatrix} 0 \\ \text{Cos}(\omega t) \\ i \text{Sin}(\omega t) \end{Bmatrix} K \sqrt{\frac{\epsilon}{\mu}} e^{-\frac{G1 \epsilon_0 \mu_0}{8 \pi r}} = K \sqrt{\frac{\epsilon}{\mu}} e^{-\frac{G1 \epsilon_0 \mu_0}{8 \pi r}} e^{i \omega t}$$


The Scalar function  $\Psi[r, t]$  represents a fundamental solution of the Quantum Mechanical Schrödinger wave equation. [36, 37]

**Black Holes with Discrete Spherical Energy Levels at Sub-Atomic Dimensions**

An essential requirement for the confinement of Electromagnetic Energy is that the Poyntingvector equals zero at the (spherical) surface of the confinement. For the confinement within a sphere, a standing electromagnetic wave pattern has been required which exists of concentric spheres, at every sphere an antinodal plane for E (or B) with a radius distance between each sphere of half the wavelength of the confinement. The constant  $k = n \pi \lambda$ , "n" a natural number (1,2,3,4.....) and  $\lambda$  the wavelength.


***Time and Radius Dependent Black Holes with Discrete Energy Levels, The Confinements of Electromagnetic Radiation within Spherical Regions:***

Every concentric sphere represents an anti-nodal surface for the Electric Field (E) or the Magnetic Field (H).



**Fig. 5: Nodal and Antinodal Spheres for Standing (Confined) Spherical Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (9)**

The Poynting Vector<sup>^</sup>:  $\bar{S} = \bar{E} \times \bar{H}$  at this spherical surface equals zero at any time and at any location at this sphere. The Electromagnetic Energy remains always within this sphere and the next concentric sphere. The concentric spheres have a difference in radius of one half wavelength of the electromagnetic radiation within the confinement and a different discrete energy level. Every concentric sphere represents an anti-nodal surface of the electric field or the magnetic field.



**Fig. 6: Nodal- and Antinodal Spheres (k = 3) for Standing (Confined) Spherical Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (9)**


Equation (24) describes a Time and Radius dependent BLACK HOLE.

$$\bar{E} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} \begin{pmatrix} 0 \\ \text{Sin}[k r] \text{Sin}[\omega t] \\ - \text{Cos}[k r] \text{Cos}[\omega t] \end{pmatrix} \quad (24)$$

$$\bar{H} = K e^{-\frac{G1\delta_0\mu_0}{8\pi r}} \sqrt{\frac{\epsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ \text{Sin}[k r] \text{Cos}[\omega t] \\ - \text{Cos}[k r] \text{Sin}[\omega t] \end{pmatrix}$$

Equation (20) represents by the function  $\text{Sin}[k r]$  ( $k = 1,2,3,4,\dots$ ) the confinement of electromagnetic radiation between two concentric spheres. K represents the amplitude of the Electric/ Magnetic Field Intensity. [14]

**Time and Polar Angle Dependent Black Holes:**




**Fig. 7: Nodal- and Antinodal Polar Angle Regions (m = 3) for Standing (Confined) Spherical Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (15)**

Equation (25) describes a Time and "Polar Angle" dependent BLACK HOLE

$$\vec{E} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} \begin{pmatrix} 0 \\ \text{Sin}[m \theta] \text{Sin}[\omega t] \\ \text{Sin}[m \theta] \text{Cos}[\omega t] \end{pmatrix} \tag{25}$$


$$\vec{H} = K e^{-\frac{G1\delta_0\mu_0}{8\pi r}} \sqrt{\frac{\epsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ \text{Sin}[m \theta] \text{Cos}[\omega t] \\ -\text{Sin}[m \theta] \text{Sin}[\omega t] \end{pmatrix}$$

Equation (19) represents by the function Sin[m θ] (m = 1,2,3,4..... ) the confinement of electromagnetic radiation between two Polar Angular Regions [15].



**Fig. 8: Nodal- and Antinodal Polar Angle Regions (m = 3) for Standing (Confined) Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (15)**

**Time and Azimuthal Angular Dependent Black Holes:**



**Fig. 9: Nodal- and Antinodal Azimuthal Angular Regions (n = 3) for Standing (Confined) Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (16)**


Equation (26) describes a Time and "Polar Angle" dependent BLACK HOLE

$$\bar{E} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} \begin{pmatrix} 0 \\ \text{Cos}[n \varphi] \text{Sin}[\omega t] \\ \text{Cos}[n \varphi] \text{Cos}[\omega t] \end{pmatrix} \quad (26)$$

$$\bar{H} = K e^{-\frac{G1\delta_0\mu_0}{8\pi r}} \sqrt{\frac{\epsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ \text{Cos}[n \varphi] \text{Cos}[\omega t] \\ -\text{Cos}[n \varphi] \text{Sin}[\omega t] \end{pmatrix}$$

Equation (26) represents by the function  $\text{Sin}[n \varphi]$  ( $n = 1,2,3,4,\dots$ ) the confinement of electromagnetic radiation between two Azimuthal Angular Regions [16].

**Time, Polar- and Azimuthal Angular Dependent Black Holes:**



**Fig. 10: Nodal- and Antinodal Polar Angular and Azimuthal Angular Regions (n = 4 and m = 4) for Standing (Confined) Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (17)**


Equation (27) describes a Time “Azimuthal Angle” and “Polar Angle” dependent BLACK HOLE

$$\bar{E} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} \begin{pmatrix} 0 \\ \text{Cos}[n \varphi] \text{Sin}[m \theta] \text{Sin}[\omega t] \\ \text{Cos}[n \varphi] \text{Sin}[m \theta] \text{Cos}[\omega t] \end{pmatrix} \quad (27)$$

$$\bar{H} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} \sqrt{\frac{\epsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ -\text{Cos}[n \varphi] \text{Sin}[m \theta] \text{Cos}[\omega t] \\ \text{Cos}[n \varphi] \text{Sin}[m \theta] \text{Sin}[\omega t] \end{pmatrix}$$

Equation (27) represents by the function  $\text{Cos}[n \varphi]$  ( $n = 1,2,3,4,\dots$ ) and  $\text{Sin}[m \theta]$  ( $m = 1,2,3,4,\dots$ ) the confinement of electromagnetic radiation between two Azimuthal Angular Regions and two Polar Angulars Regions [17].

***Spherical Confinement of Light Between Two Concentric Spheres Within Black Holes:***




**Fig.11: Nodal- and Antinodal Regions for Standing (Confined) Electromagnetic waves with a 90 degrees phase shift between the Electric field and the Magnetic field. Equation (14)**

Equation (18) represents the reflection of the Confined Electromagnetic Energy within the BLACK HOLE between two concentric spheres while the speed of light, depending on the variable "r", changes in direction with the frequency of the confined light (Electromagnetic Radiation).

A BLACK HOLE can split into two new BLACK HOLES with different radii. The original BLACK HOLE falls back into a lower energy level while the new BLACK HOLE represents the difference in Energy Levels comparable with an atom falling back into a lower energy level.

$$\bar{E} = K e^{-\frac{G1\epsilon_0\mu_0}{8\pi r}} f \left[ t - \frac{\sqrt{\epsilon_0 \mu_0} \cos[2 k r]}{2 k} \right] \begin{pmatrix} 0 \\ \sin[k r] \sin[\omega t] \\ -\cos[k r] \cos[\omega t] \end{pmatrix} \quad (28)$$

$$\bar{H} = K e^{-\frac{G1\delta_0\mu_0}{8\pi r}} f \left[ t - \frac{\sqrt{\epsilon_0 \mu_0} \cos[2 k r]}{2 k} \right] \sqrt{\frac{\epsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ -\sin[k r] \cos[\omega t] \\ -\cos[k r] \sin[\omega t] \end{pmatrix}$$



**Fig. 12: Nodal- and Antinodal Regions for Standing (Confined) Electromagnetic within two concentric spheres. Equation (18)**

### UNIVERSAL EQUILIBRIUM IN THE "CONCEPT OF QUANTUM MECHANICAL PROBABILITY" IN "THE NEW THEORY"

The 4-dimensional notation for the divergence of the Stress-Energy Tensor (25) expresses in the 4<sup>th</sup> dimension (time dimension) the law of Conservation of Energy". For an Electromagnetic Field the law for conservation of Energy has been expressed as:

$$(29) \quad \bar{f}^4 = \begin{pmatrix} f_4 \\ f_3 \\ f_2 \\ f_1 \end{pmatrix} = \square \square \bar{T} = \begin{pmatrix} \nabla \cdot \bar{S} + \frac{\partial w}{\partial t} \\ f_3 \\ f_2 \\ f_1 \end{pmatrix} = \bar{0}^4$$

From the equation for the "Conservation of Electromagnetic Energy" (38.1) the "Fundamental Equation for Confined Electromagnetic Interaction" in "The New Theory" will be derived, which equals the Relativistic Quantum Mechanical "Dirac" equation and the Schrödinger wave equation at velocities relatively low compared to the speed of light.

The “Fundamental Equation for Confined Electromagnetic Interaction” in “The Proposed Theory” can be considered to be the relativistic version of the Quantum Mechanical Schrödinger wave equation, which equals the Quantum Mechanical Dirac Equation.

### Confined Electromagnetic Energy Within a 4-Dimensional Equilibrium

The physical concept of quantum mechanical probability waves has been created during the famous 1927 5<sup>th</sup> Solvay Conference. During that period there were several circumstances which came just together and made it possible to create a unique idea of “Material Waves” (Solutions of Schrödinger’s wave equation) being complex (partly real and partly imaginary) and describing the probability of the appearance of a physical object (elementary particle) generally indicated as “Quantum Mechanical Probability Waves”.

The idea of complex (probability) waves is directly related to the concept of confined (standing) waves. Characteristic for any standing acoustical wave is the fact that the Velocity and the Pressure (Electric Field and Magnetic Field in QLT) are always shifted over 90 degrees. The same principle does exist for the standing (confined) electromagnetic waves,

For that reason every confined (standing) Electromagnetic wave can be described by a complex sum vector  $\bar{\phi}$  of the Electric Field Vector  $\bar{E}$  and the Magnetic Field Vector  $\bar{B}$  ( $\bar{E}$  has 90 degrees phase shift compared to  $\bar{B}$ ).

The vector functions  $\bar{\phi}$  and the complex conjugated vector function  $\bar{\phi}^*$  will be written as:

$$\bar{\phi} = \frac{1}{\sqrt{2\mu}} \left( \bar{B} + i \frac{\bar{E}}{c} \right) \quad (30)$$

$\bar{B}$  equals the magnetic induction,  $\bar{E}$  the electric field intensity ( $\bar{E}$  has + 90 degrees phase shift compared to  $\bar{B}$ ) and  $c$  the speed of light.

The complex conjugated vector function  $\bar{\phi}^*$  equals:

$$\bar{\phi}^* = \frac{1}{\sqrt{2\mu}} \left( \bar{B} - i \frac{\bar{E}}{c} \right) \quad (31)$$

The dot product equals the electromagnetic energy density  $w$ :

$$\bar{\phi} \cdot \bar{\phi}^* = \frac{1}{2\mu} \left( \bar{B} + i \frac{\bar{E}}{c} \right) \cdot \left( \bar{B} - i \frac{\bar{E}}{c} \right) = \frac{1}{2} \mu H^2 + \frac{1}{2} \varepsilon E^2 = w \quad (32)$$

Using Einstein’s equation  $W = m c^2$ , the dot product equals the electromagnetic mass density  $w$ :

$$\bar{\phi} \cdot \bar{\phi}^* \frac{1}{c^2} = \frac{\varepsilon}{2} \left( \bar{B} + i \frac{\bar{E}}{c} \right) \cdot \left( \bar{B} - i \frac{\bar{E}}{c} \right) = \frac{1}{2} \varepsilon \mu^2 H^2 + \frac{1}{2} \varepsilon^2 E^2 = \rho \text{ [kg/m}^3\text{]} \quad (33)$$



The cross product is proportional to the Poynting vector (Ref. 3, page 202, equation 15).

$$\bar{\phi} \times \bar{\phi}^* = \frac{1}{2\mu} \left( \bar{\mathbf{B}} + i \frac{\bar{\mathbf{E}}}{c} \right) \times \left( \bar{\mathbf{B}} - i \frac{\bar{\mathbf{E}}}{c} \right) = i \sqrt{\epsilon \mu} \bar{\mathbf{E}} \times \bar{\mathbf{H}} = i \sqrt{\epsilon \mu} \bar{\mathbf{S}} \quad (34)$$

This article presents a new “Gravitational-Electromagnetic Equation” describing Electromagnetic Field Configurations which are simultaneously the Mathematical Solutions for the Scalar Quantum Mechanical “Schrodinger Wave Equation” and more exactly the Mathematical Solutions for the Tensor representation of the “Relativistic Quantum Mechanical Dirac Equation” (41).

The 4-dimensional divergence of the sum of the Electromagnetic Stress-Energy tensor expresses the 4-dimensional Force-Density vector (expressed in [N/m<sup>3</sup>] in the 3 spatial coordinates) as the result of Electro-Magnetic-Gravitational interaction.

$$f^\mu = \partial_\nu \mathbf{T}^{\mu\nu} = 0 \quad (35)$$

In vector notation the 4-dimensional Force-Density vector can be written as:

$$\bar{f}^4 = \begin{pmatrix} f_4 \\ f_3 \\ f_2 \\ f_1 \end{pmatrix} = \square \square \bar{\mathbf{T}} = 0 \quad (36)$$

The fundamental boundary condition for this alternative approach to gravity is the requirement that the Force 4 vector equals zero in the 4 dimensions, expressing a universal 4-dimensional equilibrium: The 3 spatial components of the Force-Density vector, as a result of Electro-Magnetic-Gravitational interaction can be written as: Substituting the electromagnetic values for the electric field intensity “E” and the magnetic field intensity “H” in (36) results in the 4-dimensional representation of the Electro-Magnetic-Gravitational Fields Equation (37):

$$\begin{aligned} & \text{Energy-Time Domain} \\ (f_4) & \Leftrightarrow \nabla \cdot (\bar{\mathbf{E}} \times \bar{\mathbf{H}}) + \frac{1}{2} \frac{\partial (\epsilon_0 (\bar{\mathbf{E}} \cdot \bar{\mathbf{E}}) + \mu_0 (\bar{\mathbf{H}} \cdot \bar{\mathbf{H}}))}{\partial t} = 0 \\ & \text{3-Dimensional Space Domain} \\ \begin{pmatrix} f_3 \\ f_2 \\ f_1 \end{pmatrix} & \Leftrightarrow -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) \\ & + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) = \bar{0} \end{aligned} \quad (37)$$

In which  $f_1, f_2, f_3$ , represent the force densities in the 3 spatial dimensions and  $f_4$  represent the force density (energy flow) in the time dimension ( $4^{\text{th}}$  dimension). Equation (37) can be written as:

$$\begin{aligned} & \text{Energy-Time Domain} \\ & \text{Conservation of Energy} \\ & \text{B-7} \\ (f_4) \quad & \nabla \cdot \bar{S} + \frac{\partial w}{\partial t} = 0 \quad (38.1) \end{aligned}$$

(38)

$$\begin{aligned} & \text{3-Dimensional Space Domain} \\ & \text{B-1} \quad \text{B-2} \quad \text{B-3} \\ & \text{B-4} \quad \text{B-5} \\ \begin{pmatrix} f_3 \\ f_2 \\ f_1 \end{pmatrix} \quad & -\frac{1}{c^2} \frac{\partial (\bar{E} \times \bar{H})}{\partial t} + \epsilon_0 \bar{E} (\nabla \cdot \bar{E}) - \epsilon_0 \bar{E} \times (\nabla \times \bar{E}) + \\ & + \mu_0 \bar{H} (\nabla \cdot \bar{H}) - \mu_0 \bar{H} \times (\nabla \times \bar{H}) = \bar{0} \quad (38.2) \end{aligned}$$

The  $4^{\text{th}}$  term in equation (38.1) can be written in the terms of the Poynting vector “S” and the energy density “w” representing the electromagnetic law for the conservation of energy (Newton’s second law of motion).

**The 4-dimensional Relativistic Dirac Equation**

Substituting (32) and (34) in Equation (38.1) results in the 4-Dimensional Tensor presentation for the relativistic quantum mechanical Dirac Equation (39):

$$\begin{aligned} (x_4) \quad & \nabla \cdot (\bar{\phi} \times \bar{\phi}^*) + \frac{i}{c} \frac{\partial \bar{\phi} \cdot \bar{\phi}^*}{\partial t} = 0 \quad (39) \\ \begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} \quad & \frac{i}{c} \frac{\partial (\bar{\phi} \times \bar{\phi}^*)}{\partial t} - (\bar{\phi} \times (\nabla \times \bar{\phi}^*) + \bar{\phi}^* \times (\nabla \times \bar{\phi})) + (\bar{\phi} (\nabla \cdot \bar{\phi}^*) + \bar{\phi}^* (\nabla \cdot \bar{\phi})) = 0 \end{aligned}$$

To transform the electromagnetic vector wave function  $\bar{\phi}$  into a scalar (spinor or one-dimensional matrix representation), the Pauli spin matrices  $\sigma$  and the following matrices (Ref. 3 page 213, equation 99) are introduced:

$$\bar{\alpha} = \begin{bmatrix} 0 & \sigma \\ \sigma & 0 \end{bmatrix} \quad \text{and} \quad \bar{\beta} = \begin{bmatrix} \delta_{ab} & 0 \\ 0 & -\delta_{ab} \end{bmatrix} \quad (40)$$

The Equations (6), (32) and (34) can be written in tensor presentation as the 4-Dimensional Relativistic Quantum Mechanical Dirac Equation: [3] (Equation 102, page 213)

$$(x_4) \quad \left( \frac{i m c}{h} \bar{\beta} + \bar{\alpha} \cdot \nabla \right) \psi = - \frac{1}{c} \frac{\partial \psi}{\partial t} \quad (41.1)$$

(41)

$$\begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} \quad - \frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \quad (41.2)$$

$$+ \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \gamma_0 \bar{\mathbf{g}} (\nabla \cdot \bar{\mathbf{g}}) - \gamma_0 \bar{\mathbf{g}} \times (\nabla \times \bar{\mathbf{g}}) = \bar{0}$$

## CONCLUSIONS

Based on the assumption of the zero-rest mass of photons, General Relativity describes the interaction between Gravity and Light within a 4-dimensional curvature in Space and Time due to a gravitational field. Light follows a path defined by this curved 4-dimensional Space and Time geometry.

The new theory, describes a bi-directional separation between mass and inertia for light (photons). Inertia can only exist only in the direction of propagation of the beam of light (photons) which determines the speed of light. Mass of the beam of light (photons) can only exist in the plane perpendicular to the direction of propagation (directions of confinement), which determines the deflection of a beam of light (photons) by a gravitational field in the plane perpendicular to the direction of propagation.

BLACK HOLES (Gravitational-Electromagnetic Confinements) are fundamental solutions of the relativistic quantum mechanical Dirac equation. Black Holes represent the large impact of "Gravitational Intensity Shift" and "Gravitational RedShift" due to a gravitational field. Both phenomena can maybe observe in the future with extremely sensitive observatories at extreme low frequency levels.

The new theory describes the impact of "CURL" [38] within the gravitational fields around Black Holes and the impact on Gravitational Lensing. Gravitational "CURL" (Equation 6) is an effect which cannot be explained and calculated by General Relativity

Within a 4-Dimensional Equilibrium and taking into account the inertia- and the gravitational-force densities within the electromagnetic field configurations, Gravitational Electromagnetic Confinements (BLACK HOLES at sub-atomic dimensions) are a physical reality and are solutions of the Relativistic Quantum Mechanical Dirac Equation (39, 41) and present spherical confinements with discrete separate energy levels.

To test the proposed theory with General Relativity, an experiment [2] has been required which measures the interaction between gravity and light within a well-defined gravitational field like the gravitational field of the earth. The difference between the calculation for Gravitational RedShift, within the Gravitational Field of the Earth, in "General Relativity" and "The New Theory" is smaller than  $10^{-16}$  and cannot be determined with present observation equipment (maximum accuracy of  $10^{-15}$  for GRS). Validation of both theories requires higher accuracies.

Dark Matter does exist because of "Gravitational RedShift" and Gravitational Intensity Shift". A complete Galaxy, existing of billions bright light emitting star constellations, with a total mass of

$10^{53}$  [kg] becomes invisible for any observatory like the “James Webb Space Telescope” at the distance of  $5 \cdot 10^{21}$  [m] (which is 10 times the radius of the Milky Way Galaxy). This distance of “Gravitational Shielding” has been controlled by the mathematical solution (20) for equation (8). The gravitational field of these Galaxies has not been effected by the effect of “Gravitational RedShift” and “Gravitational IntensityShift” at all.

For this reason a large percentage of the total mass in the Universe beyond the border of “Gravitational Shielding” becomes invisible for our observatories on and close around the earth. While the gravitational fields of these galaxies still has the full influence on our universe.

Nuclear fusion represents the border area (nuclear plasma) between the material world (infusion of Deuterium) and the energy world (microwave heating). Existing theories to describe these material-energy interactions are far from the required necessary theoretical physics to realize stable nuclear fusion processes inside confinements like the Tokamak. The only possibility to describe these complex “mechanical-electromagnetic” interaction processes correctly is to develop a new theory in physics which describes the electro-magnetic-gravitational (linear- or rotational acceleration) force density interactions (expressed in  $\text{N/m}^3$ ) (equation 8) with the mechanical force density interactions (expressed in  $\text{N/m}^3$ ) being presented by the Navier-Stokes equation for compressible nuclear plasmas [41].

Equation (42) and for laminar Plasma Flows (45) represents the 3-dimensional equilibrium between all the dynamic- and electromagnetic force densities within the nuclear plasma and consequently the fundamental stability requirements for nuclear fusion.

### Data Availability

All Data and Calculations have been published at: <https://quantumlight.science/>

### REFERENCES

- [1] Wheeler; John Archibald; GEONs, Physical Review Journals Archive, 97, 511, Issue 2, pages 511-526, Published 15 January 1955, Publisher: American Physical Society, DOI: 10.1103/PhysRev.97.511:
- [2] Sven Herrmann, Felix Finke, Martin Lülfi, Olga (et. Al.) |; Test of the Gravitational Redshift with Galileo Satellites in an Eccentric Orbit; Phys. Rev. Lett. **121**, 231102 – Published 4 December 2018; Gravitational Redshift Test Using Eccentric Galileo Satellites, DOI: 10.1103/PhysRevLett.121.231102
- [3] Vegt, J. W. A Continuous Model of Matter based on AEONs, Physics Essays, Volume 8, Number 2, 1995, DOI: 10.31219/osf.io/ra7ng
- [4] Mathematical Solutions for the Propagation of Light in Quantum Light Theory, Calculations in Mathematica 13.1: [https://community.wolfram.com/groups/-/m/t/2576692?p\\_p\\_auth=mTldHX3v](https://community.wolfram.com/groups/-/m/t/2576692?p_p_auth=mTldHX3v)
- [5] Gravitational RedShift between two Atomic Clocks, Calculations in Mathematica 13.1: [https://community.wolfram.com/groups/-/m/t/2622560?p\\_p\\_auth=EC8QOoXz](https://community.wolfram.com/groups/-/m/t/2622560?p_p_auth=EC8QOoXz)
- [6] Propagation of Light within a Gravitational Field in Quantum Light Theory, Calculation in Mathematica 13.1: [https://community.wolfram.com/groups/-/m/t/2576537?p\\_p\\_auth=iljE3giH](https://community.wolfram.com/groups/-/m/t/2576537?p_p_auth=iljE3giH)
- [7] Raymond J. Beach; A classical Field Theory of Gravity and Electromagnetism; Journal of Modern Physics; 2014, 5, 928-939

- [8] Maxwell; James Clerk; A dynamical theory of the electromagnetic field; 01 January 1865; <https://royalsocietypublishing.org/doi/10.1098/rstl.1865.0008>
- [9] A. Einstein; On the Influence of Gravitation on the Propagation of Light; *Annalen der Physik (ser. 4)*, 35, 898–908, [http://myweb.rz.uni-augsburg.de/~eckern/adp/history/einstein-papers/1911\\_35\\_898-908.pdf](http://myweb.rz.uni-augsburg.de/~eckern/adp/history/einstein-papers/1911_35_898-908.pdf)
- [10] Mahendra Goray, Ramesh Naidu Annavarapu, Rest mass of photon on the surface of matter, *Results in Physics* 16 (202) 102866, January 2020, DOI: 10.1016/j.rinp.2019.102866
- [11] Genova, A., Mazarico, E., Goossens, S. et al.; Solar system expansion and strong equivalence principle as seen by the NASA MESSENGER mission; *Nat Commun* 9; 289 (2018). DOI: 10.1038/s41467-017-02558-1
- [12] John G. Williamson; A new linear theory of light and matter; 2019; *J. Phys.: Conf. Ser.* 1251 012050 DOI 10.1088/1742-6596/1251/1/012050
- [13] BLACK HOLES with Discrete Spherical Energy Levels: [https://community.wolfram.com/groups/-/m/t/2896941?p\\_p\\_auth=D7ZKu03k](https://community.wolfram.com/groups/-/m/t/2896941?p_p_auth=D7ZKu03k)
- [14] Time and Radius dependent GEONs with discrete Energy Levels. [https://community.wolfram.com/groups/-/m/t/2900869?p\\_p\\_auth=yxR9nZu6](https://community.wolfram.com/groups/-/m/t/2900869?p_p_auth=yxR9nZu6)
- [15] Time and Angular Regions dependent GEONs with discrete energy levels. [https://community.wolfram.com/groups/-/m/t/2901457?p\\_p\\_auth=H4jjDHmQ](https://community.wolfram.com/groups/-/m/t/2901457?p_p_auth=H4jjDHmQ)
- [16] Time and Azimuthal Regions dependent GEONs with discrete energy levels [https://community.wolfram.com/groups/-/m/t/2902170?p\\_p\\_auth=ytoq5nEh](https://community.wolfram.com/groups/-/m/t/2902170?p_p_auth=ytoq5nEh)
- [17] Time, Polar Angular and Azimuthal Angular Regions dependent GEONs with discrete energy levels [https://community.wolfram.com/groups/-/m/t/2902642?p\\_p\\_auth=sW2mvv9L](https://community.wolfram.com/groups/-/m/t/2902642?p_p_auth=sW2mvv9L)
- [18] D. W. Sciama; The Physical Structure of General Relativity; *Rev. Mod. Phys.* 36, 463 – Published 1 January 1964; Erratum *Rev. Mod. Phys.* 36, 1103 (1964)
- [19] Adrian del Rio, Jose Navarro-Salas, and Francisco Torrenti; Renormalized stress-energy tensor for spin -1/2 fields in expanding universes; *Phys. Rev. D* 90, 084017 – Published 13 October 2014
- [20] Stergios Pellis; Unity Formulas for the Coupling Constants and the Dimensionless Physical Constants; *Journal of High Energy Physics Gravitation and Cosmology*; DOI: 10.4236/jhepgc.2023.91021
- [21] Bloch, Yakov and Joshua Foo. "How the result of a measurement of a photon's mass can turn out to be 100." (2023). Corpus ID: 258426255
- [22] Andrés Arámburo García, Kyrylo Bondarenko, Sylvia Ploeckinger, Josef Pradler and Anastasia Sokolenko; Effective photon mass and (dark) photon conversion in the inhomogeneous Universe; *Journal of Cosmology and Astroparticle Physics*, Volume 2020, October 2020
- [23] Alexander M Gabovich and Nadezhda A Gabovich; How to explain the non-zero mass of electromagnetic radiation consisting of zero-mass photons; *European Journal of Physics*; 2007; 28 649; DOI 10.1088/0143-0807/28/4/004
- [24] Liang-Cheng Tu, Jun Luo and George T Gillies; The mass of the photon; *Reports on Progress in Physics*, Volume 68, Number 1; DOI 10.1088/0034-4885/68/1/R02
- [25] Doyon, B. Conformal Loop Ensembles and the Stress–Energy Tensor. *Lett Math Phys* 103, 233–284 (2013). <https://doi.org/10.1007/s11005-012-0594-1>

- [26] T. P. Hack and V. Moretti; On the stress–energy tensor of quantum fields in curved spacetimes—comparison of different regularization schemes and symmetry of the Hadamard/Seeley–DeWitt coefficients; 2012 J. Phys. A: Math. Theor. 45 374019; DOI: 10.1088/1751-8113/45/37/374019
- [27] Adam Levi; Renormalized stress-energy tensor for stationary black holes; Phys. Rev. D 95, 025007 – Published 10 January 2017; <https://doi.org/10.1103/PhysRevD.95.025007>
- [28] Gobbi, Julio; Luminiferous Aether: General Science Journal; December 10, 2018  $\gamma_0$  = Gravitational permeability of vacuum [kg s<sup>2</sup> m<sup>-3</sup>]
- [29] Xing-Hao Ye, Qiang Lin; Gravitational Lensing Analyzed by Graded Refractive Index of Vacuum; Journal of Optics A: Pure and Applied Optics; 1 May 2008; DOI 10.1088/1464-4258/10/7/075001
- [30] Wim Vejt; "The Origin of Gravity in "Quantum Light Theory""; OSF Preprints; October 14. doi:10.31219/osf.io/n43yd
- [31] P. Delva, N. Puchades, E. Schönemann, F. Dilssner, C. Courde (et. all); Gravitational Redshift Test Using Eccentric Galileo Satellites; Phys. Rev. Lett. 121, 231101 – Published 4 December 2018; DOI: 10.1103/PhysRevLett.121.231101
- [32] Oppenheim, Jonathan, A Postquantum Theory of Classical Gravity, Phys. Rev. X, Vol. 13, December 2023; DOI: <https://doi.org/10.1103/PhysRevX.13.041040>
- [33] Wim Vejt, The Origin of Gravity, A second order Lorentz Transformation for "Accelerated Electromagnetic Fields", Generating a Gravitational Field and the property of Mass, International Research Journal of Pure and Applied Physics Vol.9 No.1, pp.12-52, 2022.
- [34] Wim Vejt, The 4-Dimensional Dirac Equation in Relativistic Field Theory, European Journal of Applied Sciences, Vol 9, No. 1, pp 35 – 93, 2021
- [35] Wim Vejt; A Perfect Equilibrium inside a Black Hole; Wolfram Community: [https://community.wolfram.com/groups/-/m/t/3087823?p\\_p\\_auth=dpH7iBMg](https://community.wolfram.com/groups/-/m/t/3087823?p_p_auth=dpH7iBMg)
- [36] Albert Einstein, "Elementare Überlegungen zur Interpretation der Grundlagen der Quanten-Mechanik", Translated into English, 2011, DOI: <https://doi.org/10.48550/arXiv.1107.3701>
- [37] Nikko John Leo S. Lobos, Reggie C. Pantig; Generalized Extended Uncertainty Principle Black Holes: Shadow and lensing in the macro- and microscopic realms; Physics 2022, 4(4), 1318-1330; <https://doi.org/10.3390/physics4040084>
- [38] Zihua Weng, Influence of velocity curl on conservation laws, October 2008, <https://doi.org/10.48550/arXiv.0810.0065>
- [39] Peter Vadasz, Rendering the Navier–Stokes Equations for a Compressible Fluid into the Schrödinger Equation for Quantum Mechanics, MDPI-Fluids, May 2016, <https://doi.org/10.3390/fluids1020018>
- [40] Young-Sam Kwon, Asymptotic limit for rotational quantum compressible Navier–Stokes equations with multiple scales, Journal of Mathematical Analysis and Applications, Volume 464, Issue 2, 15 August 2018, Pages 1408-1424, <https://doi.org/10.1016/j.jmaa.2018.04.073>
- [41] Senjo Shimizu and Hidenobu Tsuritani, on a Navier–Stokes–Ohm problem from plasma physics in multi connected domains, Partial Differential Equations and Applications (2021) 2:75, <https://doi.org/10.1007/s42985-021-00122-7>
- [42] Jean-Luc Cambier and David A. Micheletti, Theoretical Analysis of the Electron Spiral Toroid Concept, NASA/CR-2000-210654, <https://ntrs.nasa.gov/api/citations/20010021117/downloads/20010021117.pdf>

[43] Soraya Field Fiorio, Isaac Newton: Magician, <https://parabola.org/2020/11/01/isaac-newton-magician/>

[44] Egil Asprem, *The problem of disenchantment: scientific naturalism and esoteric discourse*, Amsterdam University, 1939

[45] *The Emerald Tablet of Hermes & The Kybalion*; ISBN-13: 9781911405269

[46] J.D. Sethian, M. Friedman e.a.; Fusion energy with lasers, direct drive targets, and dry wall chambers; *Nuclear Fusion*, Volume 43, Number 12; DOI 10.1088/0029-5515/43/12/015